

# Assessing the *in vivo* effects of *Aloysia citrodora* extract: data from K14-HPV16 transgenic mice

Medeiros-Fonseca B<sup>1,2</sup>, Vala H<sup>\*1,3,4</sup>, Vasconcelos-Nóbrega C<sup>1,3</sup>, Faustino-Rocha A<sup>1,5,6</sup>, Medeiros R<sup>2</sup>, Bastos MMSM<sup>7,8</sup> da Costa RG<sup>1,2,7,8,9</sup> and Oliveira PA<sup>1</sup>

- 1- Centre for the Research and Technology of Agro-Environmental and Biological Sciences (CITAB), University of Trás-os-Montes and Alto Douro, UTAD, Vila Real, Portugal
- 2- Molecular Oncology and Viral Pathology Group, Research Center of IPO Porto, Portugal
- 3- Agrarian Superior School of Viseu, Polytechnic Institute of Viseu, Portugal.
- 4-Educational, Technologies and Health Study Center, Polytechnic Institute of Viseu, Portugal.
- 5-Department of Zootechnics, School of Sciences and Technology, University of Evora, Evora, Portugal.
- 6-Comprehensive Health Research Centre, Evora, Portugal.
- 7-Laboratory for Process Engineering, Environment, Biotechnology and Energy (LEPABE), Faculty of Engineering, University of Porto (FEUP), 4200-465 Porto, Portugal
- 8-Associate Laboratory in Chemical Engineering (ALICE), FEUP, 4200-465 Porto, Portugal.
- 9-Postgraduate Programme in Adult Health, UFMA, Brazil



## BACKGROUND

*Aloysia citrodora* is used in folk medicine in the preparation of infusions due to its antispasmodic, digestive, sedative, and antipyretic properties. It also has antioxidant and antimicrobial properties.

## PURPOSE OF THE EXPERIMENTAL

The present study evaluated the *in vivo* efficacy of aqueous extract (AE) of *Aloysia citrodora* in K14-HPV16 transgenic mice which HPV16 early genomic region and hence develop multiple-step epithelial neoplasia.




Figure 1. Appearance of wildtype (A) and K14-HPV16 (B) mice.

## METHODS

Thirty wildtype (WT) or K14-HPV16 (figure 1) female mice were divided into six groups (G) (n=5): G1 (WT, control), G2 (HPV, control), G3 (WT, 0.013g/mL), G4 (HPV, 0.006g/mL), G5 (HPV, 0.008g/mL), G6 (HPV, 0.013g/mL). AE was prepared and provided in drinking water every 48 h; the experiment lasted 28 days. Body weight, food and water consumption and humane endpoints were recorded weekly. At necropsy, chest and ear skin was collected and processed for histological analysis (figure 2).




Figure 2. Summary of the experimental design. Extract (A), administration in drinking water (B), samples collection (C), and histological analysis (D).

## RESULTS

The mean body weight of all animals increased throughout the experiment ( $p < 0.05$ ). Food and water intake were slightly higher in the HPV groups. The humane endpoints score was different between G2 and G4, and G4 and G5 ( $p < 0.05$ ) (table 1). The skin lesions in K14-HPV16 mice comprised hyperplasia, dysplasia, benign tumours (papillomas) and malignant tumours (carcinomas), with the worst lesions registered on the ear pavilion in G2, followed by G5 ( $p < 0.05$ ). The highest dose group (G6) showed a lower number of malignant tumours ( $p < 0.05$ ) (figure 3-5).

Group	Humane endpoints (M ± SD)
G1 WT water	0.08 ± 0.18
G2 HPV water	0.36 ± 0.41 <sup>1</sup>
G3 WT 0.013 g/mL	0.24 ± 0.22
G4 HPV 0.006 g/mL	0.36 ± 0.26
G5 HPV 0.008 g/mL	1.04 ± 0.09 <sup>2</sup>
G6 HPV 0.013 g/mL	0.80 ± 0.40

(1) Statistically different from group 4 ( $p < 0.05$ ); (2) Statistically different from group 4 ( $p < 0.05$ ).




Figure 3. Histological results of ear skin.




Figure 4. Histological results of chest skin.




Figure 5. Representative images of the skin lesions observed in mice under study (H&E staining). Ear skin (A) and chest skin (B). Ear skin in (G1) normal, (G2) carcinoma invasive, (G3) normal, (G4) papillomatosis, (G5) carcinoma *in situ*, (G6) papilloma. Chest skin, in (G1) normal, (G2) carcinoma *in situ*, (G3) normal, (G4) dysplasia, (G5) hyperplasia, (G6) papilloma.

## CONCLUSION

The results suggest that AE is safe for the studied animals, showing a dose-dependent trend towards lesion improvement. As AE is a natural product, the authors believe that further studies including higher doses are now warranted.

## ACKNOWLEDGMENTS

BMF is supported by FCT through grant number 2020.07675.BD. This work is supported by National Funds from FCT under the projects UIDB/04033/2020 and UIDB/00681/2020 (CERNAS9).